



Fall-Applied Herbicides: Which Weed Species Should be the Target?

Aaron Hager

Department of Crop Sciences
University of Illinois

August 22, 2018

farmdoc daily (8): 157

Recommended citation format: Hager, A. "Fall-Applied Herbicides: Which Weed Species Should be the Target?" *farmdoc daily* (8): 157, Department of Agricultural and Consumer Economics, University of Illinois at Urbana-Champaign, August 22, 2018.

Permalink: <https://farmdocdaily.illinois.edu/2018/08/fall-applied-herbicides-which-weed-species-should-be-the-target.html>

While the 2018 Illinois corn and soybean crops await harvest, assessing weed control successes and failures now can help farmers plan strategies for 2019. The practice of applying herbicides after crop harvest has increased in Illinois over the past few seasons. Herbicides applied in the fall often can provide improved control of many winter annual weed species compared with similar applications made in the spring. Horseweed (a.k.a. marestail) (Figure 1) is one example of a weed species that is often better controlled with herbicides applied in the fall compared with the spring. An increasing frequency of horseweed populations in Illinois are resistant to glyphosate, and resistance to ALS-inhibiting herbicides also is present in Illinois populations. Targeting emerged horseweed with higher application rates of products such as 2,4-D in the fall almost always results in better control at planting compared with targeting overwintered and often larger plants with lower rates of 2,4-D in the spring.


fdd

Figure 1. Young Horseweed Plant



We request all readers, electronic media and others follow our citation guidelines when re-posting articles from farmdoc daily. Guidelines are available [here](#). The farmdoc daily website falls under University of Illinois copyright and intellectual property rights. For a detailed statement, please see the University of Illinois Copyright Information and Policies [here](#).

One question typically posed is whether or not a fall application needs to include one or more herbicides that provide soil residual control of winter annual weed species. Typically, the earlier the fall application is made (say, early October) the more benefit a soil-residual herbicide can provide since emergence of winter annual weeds is often not complete. However, delaying the herbicide application until later in the fall (say, mid-November) often diminishes the necessity of a soil-residual herbicide since most of the winter annual weeds have emerged and can be controlled with non-residual herbicides. Butterweed, for example (Figure 2), is a winter annual species that research has demonstrated essentially completes its emergence by November. In University of Illinois research, fall-applied glyphosate generally provided greater and more consistent control of butterweed compared to the same rate applied in spring.



Applying a soil-residual herbicide late in the fall in hopes of having a clean field prior to planting is akin to gambling on the weather. Cold winter conditions can reduce herbicide degradation in the soil and increase herbicide persistence. This might not always be favorable since, depending on the residual herbicide, increased persistence also can cause injury to the following crop. A more moderate winter and early spring warming will increase herbicide degradation, which could result in the need for a burndown herbicide to control existing vegetation before planting.

We recommend fall-applied herbicides target fall-emerging winter annual species, biennials and perennials. We do not recommend fall application of residual herbicides for control of any spring-emerging annual weed species (especially *Amaranthus* species such as waterhemp and Palmer amaranth) the next spring for the following reasons:

- 1) Inconsistent performance: as previously described, the performance consistency of soil-residual herbicides applied in the fall is greatly dependent on weather and soil conditions after application. Our data suggest the greatest and most consistent control of *Amaranthus* species either at planting or several weeks after planting was achieved when residual herbicides were applied in the spring, not in the fall.
- 2) Increased selection for herbicide-resistant biotypes: soil-applied herbicides are not immune from selection for herbicide-resistant biotypes. Following a fall application, the concentration of herbicide remaining in the spring when *Amaranthus* species begin to germinate will be much lower compared with the same product rate applied closer to planting.

Populations of several summer annual broadleaf weed species in Illinois demonstrate resistance to herbicides from more than one site-of-action herbicide class. Their effective management requires an integrated approach that often includes soil-residual herbicides. Applying these herbicides when they will be most effective against these challenging summer annual species is a critical component of an integrated management program.